

Product Information



Sweetlix

LIVESTOCK SUPPLEMENT SYSTEM

#11386
SWEETLIX® EnProAl® 20% Poured

Features & Benefits

- Convenient, palatable source of protein, energy, minerals and vitamins for cattle consuming low quality forages
- Unique EnProAl® technology results in uniform hardness and consistency throughout
- Uniform consistency results in a known, predictable consumption rate
- Predictable consumption = predictable feed costs
- All natural protein is ideal for all classes of cattle
- High magnesium content helps protect against grass tetany
- No expensive bunks or feeders required
- For optimum forage utilization, provide a SWEETLIX® self- fed mineral supplement in conjunction with this product

Nutrient Guarantee

Crude Protein, Min	20.00%	Magnesium, Min	3.00%
Crude Fat, Min	5.00%	Copper, Min	45 ppm
Crude Fiber, Max	2.00%	Manganese, Min	170 ppm
Calcium, Min	0.75%	Selenium, Min	3.4 ppm
Calcium, Max	1.25 %	Zinc, Min	130 ppm
Phosphorus, Min	1.00%	Vitamin A, Min	50,000 IU/ lb
Salt, Min	2.25%	Vitamin D-3, Min	5,000 IU/ lb
Salt, Max	2.75%	Vitamin E, Min	5 IU/ lb



Product Ingredients

Molasses Products, Processed Grain By- Products, Hydrolyzed Feather Meal, Magnesium Oxide, Monocalcium Phosphate, Dicalcium Phosphate, Hydrolyzed Vegetable Oil, Salt, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Calcium Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

Feeding Directions

Introductory Period: Following a 7 to 10 day acclimation period, average consumption should be 1 to 2 pounds per head per day for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from loafing, feeding and watering areas. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.

Feeding Recommendations:

1. Feed 1 block per 10 to 20 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing, feeding and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum.
3. Feed blocks continuously along with a plentiful source of forage and clean, fresh water. Cattle should not run out of blocks. When a block is two- thirds consumed, provide a new block near it.
4. Cattle should consume 1 to 2 pounds per head daily based on a 1,000- pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional mineral and vitamin fortification is desired, provide an appropriate SWEETLIX® free choice mineral supplement in addition to this supplement.

CAUTION: USE AS DIRECTED. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

This Product Information Sheet is not intended to replace the Product Label.
Always read the Product Label for the most current nutrient content and feeding directions.

www.sweetlix.com
1-888-718-3493